



1. Demonstrated serologic evidence of measles, mumps or rubella antibodies or laboratory confirmation of these diseases is acceptable proof of immunity to these diseases. Serologic tests for polio are acceptable proof of immunity only if the test was performed before September 1, 2019, and all three serotypes were positive. A positive blood test for hepatitis B surface antibody is acceptable proof of immunity to hepatitis B. Demonstrated serologic evidence of varicella antibodies, laboratory confirmation of varicella disease or diagnosis by a physician, physician assistant or nurse practitioner that a child has had varicella disease is acceptable proof of immunity to varicella.

2. Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine.
(Minimum age: 6 weeks)

a. Children starting the series on time should receive a 5-dose series of DTaP vaccine at 2 months, 4 months, 6 months and at 15 through 18 months and at 4 years or older. The fourth dose may be received as early as age 12 months, provided at least 6 months have elapsed since the third dose. However, the fourth dose of DTaP need not be repeated if it was administered at least 4 months after the third dose of DTaP. The final dose in the series must be received on or after the fourth birthday and at least 6 months after the previous dose.

b. If the fourth dose of DTaP was administered at 4 years or older, and at least 6 months after dose 3, the fifth (booster) dose of DTaP vaccine is not required.

c. Children 7 years and older who are not fully immunized with the childhood DTaP vaccine series should receive Tdap vaccine as the first dose in the catch-up series; if additional doses are needed, use Td or Tdap vaccine. If the first dose was received before their first birthday, then 4 doses are required, as long as the final dose was received at 4 years or older. If the first dose was received on or after the first birthday, then 3 doses are required, as long as the final dose was received at 4 years or older.

3. Tetanus and diphtheria toxoids and acellular pertussis (Tdap) adolescent booster vaccine. (Minimum age for grades 6 through 10: 10 years; minimum age for grades 11 and 12: 7 years).

a. Students 11 years or older entering grades 6 through 12 are required to have one dose of Tdap.

b. In addition to the grade 6 through 12 requirement, Tdap may also be given as part of the catch-up series for students 7 years of age and older who are not fully immunized with the childhood DTaP series, as described above. In school year 2024-25, only doses of Tdap given at age 10 years or older will satisfy the Tdap requirement for students in grades 6 through 10; however, doses of Tdap given at age 7 years or older will satisfy the requirement for students in grades 11 and 12.

c. Students who are 10 years old in grade 6 and who -USgra 5BDC/P 22-US to have one dose of Tdap4BDC BT0 Tc 10 0 0 10 36 796.1001 Tm(3.)Tj/Span4ActualTetActivat,
- For further information, contact:

New York State Department of Health

Division of Vaccine Excellence

Room 649, Corning Tower ESP

Albany, NY 12237

(518) 473-4437

New York City Department of Health and Mental Hygiene

School Compliance Unit, Bureau of Immunization

42-09 28th Street, 5th floor

Long Island City, NY 11101

(347) 396-2433

N

D

m

/D

.

.

.

/mm

2370

04/24